

REMARKS**INTRODUCTION:**

In accordance with the foregoing, claims 1, 2, 3, 21, 22, 23, 25, 26, 27, 30, 35, 36, and 37 have been amended. No new matter is being presented, and approval and entry are respectfully requested.

Claims 1-12, 21-33 and 35-41 are under consideration. Claims 13-20 are withdrawn. Reconsideration is respectfully requested.


CHANGES TO THE SPECIFICATION AND CLAIMS:

The specification has been reviewed in response to this Office Action. Changes have been made to the specification only to place it in preferred and better U.S. form for issuance and to resolve the Examiner's objections raised in the Office Action. No new matter has been added.

It is respectfully submitted that there may be some confusion with respect to fluorene and fluorenone. Below are descriptions of both fluorene and fluorenone from <http://www.chemicaland21.com/arokorhi/specialtychem/finechem/FLUORENE.htm> and <http://www.chemicaland21.com/arokorhi/specialtychem/finechem/FLUORENONE.htm>, respectively:

FLUORENE**PRODUCT IDENTIFICATION**

CAS NO.	86-73-7
EINECS NO.	201-695-5
FORMULA	C ₁₃ H ₁₀
MOL WT.	166.22



H.S. CODE

TOXICITY

SYNONYMS 9H-Fluorene; o-Biphenylenemethane;
o-Biphenylmethane; DiPhenylenemethane; 2,2'-Methylenebiphenyl; 2,3-Benzindene;

RAW MATERIALS

CLASSIFICATION

PHYSICAL AND CHEMICAL PROPERTIES

PHYSICAL STATE	white solid
MELTING POINT	116 - 117 C
BOILING POINT	295 C
SPECIFIC GRAVITY	1.203
SOLUBILITY IN WATER	Insoluble
AUTOIGNITION	
pH	

VAPOR DENSITY
NFPA RATINGS
REFRACTIVE INDEX
FLASH POINT

151 C

STABILITY

Stable under ordinary conditions. Oxidizes in light

GENERAL DESCRIPTION AND APPLICATIONS

Polycyclic aromatic hydrocarbons (also called polynuclear hydrocarbons) have two or more single or fused aromatic rings if a pair of carbon atoms is shared between rings in their molecules. In particular, the term 'PAH' refers to the compounds consisting of only carbon and hydrogen atoms while the wider term 'polycyclic aromatic compounds' includes the alkyl-substituted derivatives and functional derivatives such as nitro- and hydroxy-PAH as well as the heterocyclic analogues, which contain one or more hetero atoms in the aromatic structure. PAHs exist in various combinations that manifest various functions such as light sensitivity, heat resistance, conductivity, emittability, corrosion resistance and physiological action. The simplest examples are naphthalene having two benzene rings side by side and biphenyl having two bond-connected benzene rings. PAHs are not found in synthetic products and are non-essential for the growth of living cells. The general characteristics of PAH describe high melting- and boiling-points (they are solid), low vapour pressure, and very low water solubility, decreasing with increasing molecular weight whereas resistance to oxidation, reduction, and vapourization increases. Vapour pressure tends to decrease with increasing molecular weight. PAHs are highly lipophilic and readily soluble in organic solvents. The lower molecular weight PAHs of 2 or 3 ring groups such as naphthalenes, fluorenes, phenanthrenes, and anthracenes have toxicity which tends to decrease with increasing molecular weight. PAHs are not synthesized chemically for industrial purposes but are isolated from concentrated coal-tar products (or from pyrolysis of coal hydrocarbons) followed by subsequent purification through repeated distillation and crystallization. Some PAHs such as naphthalene are also obtained from the concentration of the high boiling residual oil (and asphalt) derived from crude petroleum refinery processing. These PAHs are mostly used as intermediaries in pharmaceuticals, agricultural products, photographic products, thermosetting plastics, lubricating materials, and other chemical industries. General uses are;

- Acenaphthene: Intermediate for naphthalic acids, naphthalic anhydride (intermediate for pigments) and for acenaphthylene (intermediate for resins); Intermediate for dyes, soaps, pigments, pharmaceuticals, insecticide, fungicide, herbicide and plant growth hormones. It is used to manufacture plastics and as an agent for inducing polyploidy.
- Acridine: Dye and pharmaceutical manufacturing
- Anthracene: Its oxidation yields anthraquinone, the parent substance of a large class of dyes and pigments; diluent for wood preservatives; scintillant (for detection of high-energy radiation)
- Fluoranthene: manufacturing fluorescent and vat dyes, pharmaceuticals and agrochemicals.
- Fluorene: basic substance for production of dyes, pigments, pesticides, thermosetting plastic and pharmaceuticals; manufacturing fluorenone (mild oxidizing agent)
- Naphthalene: In the production of phthalic anhydride, carbaryl insecticide, beta-naphthol, tanning agents, moth repellent, and surfactants - naphthalene: main use: production of phthalic anhydride (intermediate for polyvinyl chloride plasticizers); also, production of azo dyes, surfactants and dispersants, tanning agents, carbaryl (insecticide), alkylnaphthalene solvents (for carbonless copy paper), and use without processing as a fumigant (moth repellent)
- Phenanthrene: manufacturing phenanthrenequinone (intermediate for pesticides);

manufacturing diphenic acid (intermediate for resins)

- Pyrene: manufacturing perinon pigments
- Quinoline: solvent for resins & terpiners; decarboxylation agent; parent compound to make drugs, fungicides, biocides, alkaloids, dyes, rubber chemicals and flavoring agents

Precise PAHs, specific refined products are used also in the field of electronics, functional plastics and liquid crystals. Pharmaceutical and agricultural PAHs obtained coal tar are such materials as indole, indolizine, indene, quinoline, quinaldine, isoquinoline and their derivatives. High boiling-point special solvent are such materials as tetralin, decaline, methyl-naphthalenes. Coumarins and dihydrocoumarins which can be obtained coal tar are PAHs used in perfumery.

Thermosensitive paper sensitizer PAHs are such materials as p-benzylbiphenyl and ethylbiphenyl. Fluorene is a member of polycyclic aromatic hydrocarbon (PAH). Two benzene rings are fused to cyclopentane ring. It emits violet fluorescent color. It is not synthesized commercially but is obtained from middle oil fraction of coal tar. It is insoluble in water; soluble in ether and acetone; melting point 116-117 C. It plays important part in metallocene catalysts as a ligand. It is used in the formation of polyradicals for resins. It is used in manufacturing antimalaria drugs and other pharmaceuticals. Fluorene family compounds are base materials for dyes and optical brightening agents. They have useful functions such as light and temperature sensitivity, heat resistance, conductivity, emittability, corrosion resistance and detection of amino groups. They are used in the applications of thermo and light sensitizer, liquid crystal chemistry, luminescence chemistry, spectrophotometric analysis, molecular chemistry, organometallic-complexes and biochemorphy industry.


SALES SPECIFICATION

APPEARANCE	white flakes
ASSAY (G.C)	97.0% min
MELTING POINT	115 ± 1 C
LOSS ON DRYING	0.5% max
TRANSMITTANCE	99.0% (50nm, 10% Toluene Sol.)


TRANSPORTATION

PACKING	25kgs in fiber drum
HAZARD CLASS	3
UN NO.	1224


EXAMPLES OF PARENT PAH COMPOUNDS




PENTALENE




INDENE (CAS RN: 95-13-6)




NAPHTHALENE (CAS RN: 91-20-3)




AZULENE (CAS RN: 275-51-4)




HEPTALENE




BIPHENYLENE (CAS RN: 259-79-0)




α S-INDACENE




s-INDACENE




ACENAPHTHALENE (CAS RN: 83-32-9)




FLUORENE (CAS RN: 86-73-7)




PHENALENE (CAS RN: 203-80-5)




ANTHRACENE (CAS RN: 120-12-7)




FLUORANTHENE (CAS RN: 206-44-0)




ACEPHENANTHRYLENE




TRIPHENYLENE (CAS RN: 217-59-4)




PYRENE (CAS RN: 129-00-0)




CHRYSENE (CAS RN: 218-01-9)




NAPHTHACENE (CAS RN: 92-24-0)




PLEIADENE (CAS RN:)




PICENE (CAS RN: 213-46-7)




PERYLENE (CAS RN: 198-55-0)




PENTAPHENE (CAS RN: 222-93-5)




PENTACENE (CAS RN: 135-48-8)




TETRAPHENYLENE (CAS RN: 212-74-8)




RUBICENE (CAS RN: 197-61-5)



CORONENE (CAS RN: 191-07-1)



PYRANTHRENE (CAS RN: 191-13-9)




OVALENE (CAS RN: 190-26-1)

9-FLUORENONE

PRODUCT IDENTIFICATION

CAS NO. 486-25-9
 EINECS NO. 207-630-7
 FORMULA $C_{13}H_8O$
 MOL WT. 180.21



H.S. CODE

TOXICITY

SYNONYMS Fluoren-9-one; 9-Fluorenone; 9H-Fluoren-9-one;
 9-Oxofluorene; Diphenylene ketone;

DERIVATION

CLASSIFICATION

PHYSICAL AND CHEMICAL PROPERTIES

PHYSICAL STATE yellow crystalline powder

MELTING POINT 82 - 85 C

BOILING POINT 342 C

SPECIFIC GRAVITY 1.13

SOLUBILITY IN WATER

AUTOIGNITION

pH

VAPOR DENSITY

NFPA RATINGS

REFRACTIVE INDEX

FLASH POINT 163 C

STABILITY Stable under ordinary conditions. Oxidizes in light

APPLICATIONS

Fluorene is a member of polycyclic aromatic hydrocarbon (PAH). Two benzene rings are fused to cyclopentane ring. It emits violet fluorescent color. It is not synthesized commercially but is obtained from middle oil fraction of coal tar. It is insoluble in water; soluble in ether and acetone; melting point 116-117 C. It plays important part in metallocene catalysts as a ligand. It is used in the formation of polyradicals for resins. It is used in manufacturing antimalaria drugs and other pharmaceuticals. Fluorene family compounds are base materials for dyes and optical brightening agents. They have useful functions such as light and temperature sensitivity, heat resistance, conductivity, emittability, corrosion resistance and detection of amino groups. They are used in the applications of thermo and light sensitizer, liquid crystal chemistry, luminescence chemistry, spectrophotometric analysis, molecular chemistry, organometallic-complexes and biochemorphology industry. Oxidation of fluorenean occur readily at the most reactive position 9. 9-Fluorenone has been investigated as an attractive element in organic solar cells, and display devices.

SALES SPECIFICATION

APPEARANCE yellow crystalline powder

ASSAY (G.C) 99.5% min

MELTING POINT 82 C min

LOSS ON DRYING 0.5% max

TRANSPORTATION

PACKING 25kgs in fiber drum

HAZARD CLASS

UN NO.


GENERAL DESCRIPTION OF PAHs

Polycyclic aromatic hydrocarbons (also called polynuclear hydrocarbons) have two or more single or fused aromatic rings if a pair of carbon atoms is shared between rings in their molecules. In particular, the term 'PAH' refers to the compounds consisting of only carbon and hydrogen atoms while the wider term 'polycyclic aromatic compounds' includes the alkyl-substituted derivatives and functional derivatives such as nitro- and hydroxy-PAH as well as the heterocyclic analogues, which contain one or more hetero atoms in the aromatic structure. PAHs exist in various combinations that manifest various functions such as light sensitivity, heat resistance, conductivity, emittability, corrosion resistance and physiological action. The simplest examples are naphthalene having two benzene rings side by side and biphenyl having two bond-connected benzene rings. PAHs are not found in synthetic products and are non-essential for the growth of living cells. The general characteristics of PAH describe high melting- and boiling-points (they are solid), low vapour pressure, and very low water solubility, decreasing with increasing molecular weight whereas resistance to oxidation, reduction, and vapourization increases. Vapour pressure tends to decrease with increasing molecular weight. PAHs are highly lipophilic and readily soluble in organic solvents. The lower molecular weight PAHs of 2 or 3 ring groups such as naphthalenes, fluorenes, phenanthrenes, and anthracenes have toxicity which tends to decrease with increasing molecular weight. PAHs are not synthesized chemically for industrial purposes but are isolated from concentrated coal-tar products (or from pyrolysis of coal hydrocarbons) followed by subsequent purification through repeated distillation and crystallization. Some PAHs such as naphthalene are also obtained from the concentration of the high boiling residual oil (and asphalt) derived from crude petroleum refinery processing. These PAHs are mostly used as intermediaries in pharmaceuticals, agricultural products, photographic products, thermosetting plastics, lubricating materials, and other chemical industries. General uses are;


- Acenaphthene: Intermediate for naphthalic acids, naphthalic anhydride (intermediate for pigments) and for acenaphthylene (intermediate for resins); Intermediate for dyes, soaps, pigments, pharmaceuticals, insecticide, fungicide, herbicide and plant growth hormones. It is used to manufacture plastics and as an agent for inducing polyploidy.
- Acridine: Dye and pharmaceutical manufacturing
- Anthracene: Its oxidation yields anthraquinone, the parent substance of a large class of dyes and pigments; .diluent for wood preservatives; scintillant (for detection of high-energy radiation)
- Fluoranthene: manufacturing fluorescent and vat dyes, pharmaceuticals and agrochemicals.
- Fluorene: basic substance for production of dyes, pigments, pesticides, thermoset plastic and pharmaceuticals; manufacturing fluorenone (mild oxidizing agent)
- Naphthalene: In the production of phthalic anhydride, carbaryl insecticide, beta-naphthol, tanning agents, moth repellent, and surfactants - naphthalene: main use: production of phthalic anhydride (intermediate for polyvinyl chloride plasticizers); also, production of azo dyes, surfactants and dispersants, tanning agents, carbaryl (insecticide), alkylnaphthalene solvents (for carbonless copy paper), and use without processing as a fumigant (moth repellent)
- Phenanthrene: manufacturing phenanthrenequinone (intermediate for pesticides); manufacturing diphenic acid (intermediate for resins)
- Pyrene: manufacturing perinon pigments
- Quinoline: solvent for resins & terpenes; decarboxylation agent; parent compound to make drugs, fungicides, biocides, alkaloids, dyes, rubber chemicals and flavoring agents

Precise PAHs, specific refined products are used also in the field of electronics, functional plastics and liquid crystals. Pharmaceutical and agricultural PAHs obtained from coal tar are such materials as indole, indolizine, indene, quinoline, quinaldine, isoquinoline and their derivatives. High boiling-point special solvent are such materials as tetralin, decaline, methyl-naphthalenes. Coumarins and dihydrocoumarins which can be obtained from coal tar are PAHs used in perfumery. Thermosensitive paper sensitizer PAHs are such materials as p-benzylbiphenyl and ethylbiphenyl.


EXAMPLES OF PAH PARENT COMPOUNDS




PENTALENE




INDENE (CAS RN: 95-13-6)




NAPHTHALENE (CAS RN: 91-20-3)




AZULENE (CAS RN: 275-51-4)




HEPTALENE




BIPHENYLENE (CAS RN: 259-79-0)




α-INDACENE




s-INDACENE




ACENAPHTHALENE (CAS RN: 83-32-9)




FLUORENE (CAS RN: 86-73-7)




PHENYLENE (CAS RN: 203-80-5)




ANTHRACENE (CAS RN:120-12-7)




FLUORANTHENE (CAS RN: 206-44-0)




ACEPHENANTHRYLENE




TRIPHENYLENE (CAS RN: 217-59-4)




PYRENE (CAS RN: 129-00-0)




CHRYSENE (CAS RN: 218-01-9)




NAPHTHACENE (CAS RN: 92-24-0)




PLEIADENE (CAS RN:)




PICENE (CAS RN: 213-46-7)




PERYLENE (CAS RN: 198-55-0)




PENTAPHENE (CAS RN: 222-93-5)




PENTACENE (CAS RN: 135-48-8)




TETRAPHENYLENE (CAS RN: 212-74-8)




RUBICENE (CAS RN: 197-61-5)



CORONENE (CAS RN: 191-07-1)



PYRANTHRENE (CAS RN: 191-13-9)



OVALENE (CAS RN:190-26-1)

It is respectfully submitted that in <http://www.chem.wisc.edu/~newtrad/CurrRef/Glossary/Glossary.html>, a derivative is defined as follows:

Derivative

A derivative of a compound resembles the original compound, except that some modifications in atomic structure are evident. Usually, derivatization of a molecule involves altering part of it slightly or adding a new part to the original compound.

Hence, Formula 1 and Formula 2 of the present claimed invention are clearly derivatives of fluorene. Also, Formula 1 and Formula 2 of the present claimed invention are derivatives of 9-fluorenone (also referred to as fluorenone). Thus, Formula 1 and Formula 2 of the present claimed invention may be described by those skilled in the art as being a fluorene compound or a fluorenone compound.

However, it appears that the Examiner believes that it may be more clear if the compounds of Formula 1 and Formula 2 of the present invention are referred to as compounds of 9-fluorenone. Hence, the terminology "fluorene" has been amended to recite ---9-fluorenone---. This change has been made throughout the specification and the claims. It is respectfully submitted that this is simply alternate naming, and it is clear to one skilled in the art that no new matter is added.

REJECTION UNDER 35 U.S.C. §112:

In the Office Action, at page 2, numbered paragraph 2, claims 1-12, 21-33 and 35-41 were rejected under 35 U.S.C. §112, first paragraph, for the reasons set forth therein. This rejection is traversed and reconsideration is requested.

The Examiner submits: "The claim(s) contains subject matter, which was not described in the specification in such a way as to reasonably convey to one skilled in the relevant art that the inventor(s), at the time the application was filed, had possession of the claimed invention. Applicants fail to teach the fluorenone compound or how to make or use the compound. The halogen (X) is not taught in the specification and there are no working examples of the halogen substitutive fluorenone. The A and B substituted are not taught on any example and the carboxyl fluorenone of claims 2, 14, 22, 26, 36 fail to have the A group if the B group is the carbonyl group."

As noted above, the specification and claims have been reviewed, and the terminology "fluorene" has been amended to recite ---9-fluorenone---. This change has been made throughout the specification and the claims for clarity.

In response to the Examiner's request for examples of how to make or use the fluorenone compound, the following examples, which are known to those skilled in the art, are provided:

Synthesis Example 1: 9-fluorenone-4-carboxylic acid

To a 1-liter 3-neck round bottom flask, equipped with a thermometer, a mechanical stirrer and a reflux condenser were added 460 g of concentrated sulfuric acid (4.7 moles, analytical grade, commercially obtained from Sigma-Aldrich, Milwaukee, WI) and 100 g of diphenic acid (0.41 mole, commercially obtained from Acros Fisher Scientific Company Inc., Hanover Park, IL). Using a heating mantle, the flask was heated to 135-145°C for 12 minutes, and then cooled to room temperature (RT). After being cooled to RT, the solution was added to a 4 liter Erlenmeyer flask containing 3 liters of water. The mixture was stirred mechanically and was boiled gently for one hour. A yellow solid was filtered out hot, washed with hot water until the pH of the washing water was neutral, and was dried in the air overnight. The yellow solid was fluorenone-4-carboxylic acid (75 g., 80% yield, m.p. 223-224°C). A ¹H-NMR spectrum of fluorenone-4-carboxylic acid was obtained in d₆-DMSO by a 300 MHz NMR from Bruker Instrument. The peaks were found at δ = 7.39-7.50 (m, 2H); δ = 7.79-7.70 (q, 2H); δ = 7.74-7.85 (d, 1H); δ = 7.88-8.00 (d, 1H); and δ = 8.18-8.30 (d, 1H), where d is doublet, t is triplet, m is multiplet, dd is double doublet, and q is quintet.

Synthesis Example 2: n-Butyl 9-fluorenone-4-carboxylic ester

To a 2-liter round bottom flask equipped with a mechanical stirrer and a reflux condenser with a Dean Stark apparatus were added 70 g (0.312 mole) of fluorenone-4-carboxylic acid, 480 g (6.5 mole) of n-Butanol (commercially obtained from Fisher Scientific Company Inc., Hanover Park, IL), 1000 ml of Toluene and 4 ml of concentrated sulfuric acid. The solution was refluxed


for 5 hours with aggressive agitation and refluxing, during which ~6 g of water were collected in the Dean Stark apparatus. The flask was cooled to room temperature. The solvents were evaporated and the residue was added to 4-liters of 3% sodium bicarbonate aqueous solution with agitation. The solid was filtered off, washed with water until the pH of the water was neutral, and dried in the hood overnight. The product was n-butyl 9-fluorenone-4-carboxylate ester (70 g, 80% yield). A ^1H -NMR spectrum of n-butyl fluorenone-4-carboxylate ester was obtained in CDCl_3 by a 300 MHz NMR from Bruker Instrument. The peaks were found at $\delta = 0.87\text{--}1.09$ (t, 3H); $\delta = 1.42\text{--}1.70$ (m, 2H); $\delta = 1.75\text{--}1.88$ (q, 2H); $\delta = 4.26\text{--}4.64$ (t, 2H); $\delta = 7.29\text{--}7.45$ (m, 2H); $\delta = 7.46\text{--}7.58$ (m, 1H); $\delta = 7.60\text{--}7.68$ (dd, 1H); $\delta = 7.75\text{--}7.82$ (dd, 1H); $\delta = 7.90\text{--}8.00$ (dd, 1H); $\delta = 8.25\text{--}8.35$ (dd, 1H); where d is doublet, t is triplet, m is multiplet, dd is double doublet, and q is quintet.

It is respectfully submitted that, in view of the above-cited amendments, the fluorenone compound and how to make or use the compound are known to those skilled in the art. For example, when $m=n=0$, one of the embodiments of Formula 1 of claim 1 is 9-fluorenone-2,7-dicarboxylic acid, which is available from Oakwood Products, Inc., 1741 Old Dunbar Road, West Columbia, South Carolina (see enclosure) and is also listed on the <http://www.chemexper.com/chemicals/supplier> listing on the Internet (see enclosure). It is respectfully submitted that it is known to those skilled in the art that halogen substituents (F, Cl, Br, I) may be added to the six-membered rings. The halogen substituents are highly electronegative, pulling electrons away from the six-membered rings toward themselves, making the rings less nucleophilic.

It is respectfully submitted that halogen is recited in amended paragraph [0024] of the specification, recited below for the convenience of the Examiner:

[0024] A composition to form the charge generating layer has a 9-fluorenone compound represented by Formula 1 as an electron transport material and charges generated by a laser beam are easily injected into a charge transport layer and an overcoat layer:

Formula 1



wherein A and B are independently selected from the group consisting of a carboxyl group

(-COOH), a substituted or unsubstituted $\text{C}_2\text{--C}_{10}$ alkoxy carbonyl group and a substituted or unsubstituted $\text{C}_2\text{--C}_{10}$ alkyl aminocarbonyl group, **X_1 and X_2 are independently a halogen atom**, and m and n are independently an integer from 0 to 3. (emphasis added)

Hence, the halogen (X) is taught in the specification. Although specific working examples of halogen-substituted fluorenones are not provided, it is respectfully submitted that halogen-substituted fluorenones are known to those skilled in the art, e.g., 2,7,-dibromo-9-fluorenone (see <http://www.chemicaland21.com/arokorhi/specialtychem/NH/2,7-DIBROMO-9-FLUORENONE.htm>, a copy of which is enclosed herewith).

In addition, as is known to those skilled in the art, when m and/or n are >0, halogen atom(s) may be coupled to the six-membered rings to form other embodiments of Formula 1 of claim 1.

Claim 2 has been amended to independent form. Claim 2 is based on amended paragraph [0026] of the specification. Hence, the Examiner's concerns about claim 2 are now submitted to be moot.

In view of the amendment of the terminology "fluorene" to "9-fluorenone" and the amendment to claim 2, it is respectfully submitted that independent claims 1, 21, 25, 30, and 35 are in allowable form under 35 U.S.C. §112, first paragraph. Since claims 3-12, 22-24, 26-28, 31-33, and 36-41 depend from amended claims 1, 21, 25, 30 and 35, respectively, claims 3-12, 22-24, 26-28, 31-33, and 36-41 are submitted to be in allowable form under 35 U.S.C. §112, first paragraph, for at least the reasons that amended claims 1, 21, 25, 30 and 35 are in allowable form under 35 U.S.C. §112, first paragraph.

CONCLUSION:

In accordance with the foregoing, it is respectfully submitted that all outstanding objections and rejections have been overcome and/or rendered moot, and further, that all pending claims patentably distinguish over the prior art. Thus, there being no further outstanding objections or rejections, the application is submitted as being in condition for allowance which action is earnestly solicited.

If the Examiner has any remaining issues to be addressed, it is believed that prosecution can be expedited by the Examiner contacting the undersigned attorney for a telephone interview to discuss resolution of such issues.

If there are any underpayments or overpayments of fees associated with the filing of this Amendment, please charge and/or credit the same to our Deposit Account No. 19-3935.

Respectfully submitted,

STAAS & HALSEY LLP

Date: May 23, 2006 By: Darleen J. Stockley
Darleen J. Stockley
Registration No. 34,257

1201 New York Avenue, N.W.
Suite 700
Washington, D.C. 20005
Telephone: (202) 434-1500
Facsimile: (202) 434-1501




Catalog # 002569

Name: 9-Fluorenone-2,7-dicarboxylic acid

CAS:	792-26-7
Purity:	tech
MF:	C ₁₅ H ₈ O ₅
BP:	
MP:	407°
Density:	
ND20:	
RTECS:	

Price	Quantity
\$26	1g
\$115	5g



[Home](#) [Catalog Search](#) [Featured Products](#) [New Items](#) [Custom Synthesis](#) [Profile Chemistry](#) [Pilot Plant](#) [Fluorochem USA](#) [Contact Page](#) [Custom Search](#) [Daikin](#)

Oakwood Products, Inc. 1741 Old Dunbar Road, West Columbia, SC. 29172

phone: 800-467-3386 phone: 803-739-8800 fax: 803-739-6957

Enter a name, molecular formula or cas number

>>more info

Find chemicals. 792-26-7

Search

Search by structure
Advanced Search

9-Fluorenone-2,7-dicarboxylic acid


RN: 792-26-7

9-Fluorenone-2,7-dicarboxylic acid, tech.

MF: C₁₅H₈O₅


9-Fluorenone-2,7-dicarboxylic acid tech

MW: 268.22552



Click on a product name to get more information on that compound, on a supplier name to get more information on that supplier.

Supplier	Description	References & Quantities		
chempur	9-Fluorenone-2,7-dicarboxylic acid, tech.	F1260.5	5g	Get offer
chempur	9-Fluorenone-2,7-dicarboxylic acid, tech.	F1260.25	25g	Get offer
chempur	9-Fluorenone-2,7-dicarboxylic acid tech	f12569.1	1g	Get offer
chempur	9-Fluorenone-2,7-dicarboxylic acid tech	f12569.5	5g	Get offer
chemos	9-Fluorenone-2,7-dicarboxylic acid	on request		Get offer
abcr	9-Fluorenone-2,7-dicarboxylic acid , 98%	OA2569	1 g	Get offer
		OA2569	5 g	Get offer
matrixscientific		004908	1g	Get offer
		004908	5g	Get offer
fluorochem	9-Fluorenone-2,7-dicarboxylic acid , tech	002569	1g	Get offer
		002569	5g	Get offer
oakwood	9-Fluorenone-2,7-dicarboxylic acid , tech	002569	1g	Get offer
		002569	5g	Get offer



Results 1-9 of 9

Local NursesJobs available in the Richmond area RN's,
LPN's, and CNA's
www.healthstaffinc.com**Entry**Free Legal Resource. Entry
OnInformation.com**Free IBM J2EE App Server**Based on open src Apache Geronimo. Comes
with free Eclipse plug-in.
www.ibm.com/developerworks/[Ads by Google](#)[Advertise on this site](#)

List of top chemicals. Visit our website to find more information like suppliers, MSDS, infra-red (IR), nuclear magnetic resonance spectra (NMR), bp, mp, nd20, molecular formula (MF), molfile, sdf file, structure, 3d model. You will also find information like safety, risk, hazard and MSDS. This database is a catalog of over 300000 chemicals and over 350 chemical suppliers. If you are a supplier you may also add your own catalog of chemicals for free.

Iupac name	RN (CAS)
5-Bromo-6-chloro-3-indolyl phosphate p-toluidine salt	6769-80-8
Indium(III) iodide	13510-35-5
3,4,5,6-Tetrabromo-o-cresol	576-55-6
5,6-Diamino-1,3-Dimethyluracil	5440-00-6
Fmoc-beta-cyclobutyl-D-Alanine	478183-63-0
Fmoc-beta-cyclobutyl-L-alanine	478183-62-9
2-Chloro-6-fluorophenylacetonitrile	75279-55-9
Mercury thiocyanatocobaltate(II)	27685-51-4
Cobalt tetrathiocyanatomercurate(II)	
9-Fluorenone-2,7-dicarboxylic acid	792-26-7
5-Fluoroindole-3-carboxaldehyde	2338-71-8
2-Methoxy-N-methylaniline	10541-78-3
Benzo[b]thiophene-2-carboxylic hydrazide	175135-07-6
Thianaphthene-2-carboxylic hydrazide	
4-Chlorobutyl acetate	6962-92-1
2,4,6-Trichlorobenzoic acid	50-43-1
1H,6H-Perfluorohexane	336-07-2
3-(5-Chloro-2-hydroxyphenyl)pyrazole	18704-67-1
4-(Trifluoromethylthio)benzaldehyde	4021-50-5
2-Mercapto-4-methylpyrimidine hydrochloride	6959-66-6
4-Methyl-2-pyrimidinethiol hydrochloride	
1H-Benzimidazole-2-methanol	4856-97-7
1,2,3-Thiadiazole-4-carboxylic acid	4100-13-4
4-Carboxy-1,2,3-thiadiazole	
Midazolam HCL	59467-96-8
3-Iodophenylboronic acid	221037-98-5
3-Hydroxy-2-quinoxalinecarboxylic acid	1204-75-7
N,N-Dimethyl-N'-ethylethylenediamine	123-83-1
(+)-3-Methyladipic acid	623-82-5
(+)-3-Methylhexanedioic acid	
Methylhexanedioic acid	
1-Nitro-2-Naphthol	550-60-7
3-Cyanobenzok Acid Methyl Ester	13531-48-1
2-(4-Biphenyl)ethylamine	17027-51-9
4,5-Dibromothiophene-2-carboxaldehyde	38071-22-6
4,5-Dibromo-thiophene-2-carbaldehyde	

3-Chloro-3-(4-fluorophenyl)acrylonitrile	126417-76-3
4,5-Dibromo-o-xylene	24932-48-7
Ethyl 2-chloro-4-(trifluoromethyl)pyrimidine-	187035-79-6
6-Bromo-2-naphthyl-beta-D-galactopyranoside	15572-30-2
1,2-Bis(2-aminophenoxy)-ethane-N,N,N',N'-tetraacetic acid tetrasodium salt BAPTA tetrasodium salt	126824-24-6
1,3-Diphenylbenzo[c]thiophene	16587-39-6
1,3-Diphenyl isothianaphthene	
4-Methoxybenzyl isocyanate	56651-60-6
cis-2-(4-Fluorobenzoyl)-1-cyclohexane-carboxylic acid	154810-33-0
5-Chloro-2-hydroxy-4-methylbenzophenone	68751-90-6
1-Acetoxy-4-diethylamino-2-butyne	22396-77-6
4-(tert-Butoxycarbonylaminomethyl)piperidine	135632-53-0
Ammonium tetrachloroplatinate(II)	13820-41-2
3,5-Dibromo-4-hydroxyacetophenone	2887-72-1
2-Methyl-5-phenylbenzoxazole	61931-68-8
Pentafluoropropionic anhydride	356-42-3
2-Bromo-4-nitrotoluene	7745-93-9
2-Methylsulphonylbenzoic acid	33963-55-2
2-(Methylsulfonyl)benzoic acid	
2-Cyanoethylhydrazine	353-07-1
Trifluoroacetic acid 2,2,2-trifluoroethyl ester	407-38-5
2,2,2-Trifluoroethyl trifluoroacetate	
Potassium tris(1-pyrazolyl)borohydride	18583-60-3
5-Fluorosalicylic acid	345-16-4
4-Fluoro-3-nitrotoluene	446-11-7
Tetradecylbenzene	1459-10-5
1-Phenyltetradecane	
Myristylbenzene	
Sultamicillin Tosylate (Patented-No Supply)	83105-70-8
4-Fluorobenzyl triphenyl phosphoniumchloride	3462-95-1
1,3,6-Naphthalenetrisulfonic acid, sodium salt, hydrate	19437-42-4
3,5-Dimethoxybenzoic Acid Methyl Ester	2150-37-0
Doyle dirhodium catalyst - RH2(4S-MEOX)4	167693-36-9
Thioacetanilide	637-53-6
3-Aminophenyl trifluoromethyl sulfone	426-59-5
1-Bromo-3,5-dichloro-2,4,6-trifluorobenzene	24812-13-3
3-Bromo-4-methoxytoluene	22002-45-5
7-amino-3-((1h-1,2,3-triazol-5-yl)-thiomethyl)-ceph-3-em-4-carboxylic acid	37539-03-0
methyl 4-(cyanomethyl)benzoate	76469-88-0
2,3,4,5,6-Pentafluorobenzophenone	1536-23-8
1,2-Dichlorohexafluorocyclobutane	356-18-3

Phenylbutazone Sodium Salt	129-18-0
1,2,3,6,7,8,11,12-Octahydrobenzo[e]pyren-9(10H)-one	68151-08-6
9-Oxo-1,2,3,6,7,8,9,10,11,12-decahydrobenzo[e]pyrene	
3,3,4,5,5,5-Hexafluoropentan-2-ol	2711-81-1
cis,cis,cis,cis-1,2,3,4-Cyclopentanetetracarboxylic acid	3786-91-2
(S)-2-(1-Naphthylmethyl)succinic acid-1-methyl ester	130693-96-8
(S)-3-Methoxycarbonyl-4-(1-naphthyl)butyric acid	
N-BOC-L-Aspartic acid	13726-67-5
N-(tert-Butoxycarbonyl)-L-aspartic acid	
Boc-L-Asp-OH	
BOC-L-Aspartic Acid-OH	
4-Chloro-3-nitrobenzonitrile	939-80-0
Nefazodone	83366-66-9
1-Fluoropyridinium tetrafluoroborate	107264-09-5
Hexachlorodisiloxane	14986-21-1
4-(4-Methylphenyl)butyric acid	4521-22-6
4-(p-Tolyl)butyric acid	
R-(-)-2-Oxothiazolidine-4-Carboxylic Acid	19771-63-2
Tetraethyl pyrophosphite	21646-99-1
Acetoniltriphenylphosphonium chloride	1235-21-8
Acetonyl triphenylphosphonium chloride	
CIS-1,4-Cyclohexanediol	931-71-5
Benzofurazan-5-carbonyl chloride	126147-86-2
perfluorotripropylamine	338-83-0
4-Difluoromethoxy-3-hydroxybenzaldehyde	151103-08-1
3,5-Dichlorobenzenesulfonyl chloride	705-21-5
3-Perfluorooctyl-1,2-epoxypropane	38565-53-6
Tetrafluorosuccinyl difluoride	679-13-0
1,4-Cyclohexanedione mono-2,2-dimethyltrimethylene ketal	69225-59-8
3,3-Dimethyl-1,5-dioxaspiro[5.5]undecan-9-one	
4-(Chloromethyl)-3,5-dimethylisoxazole	19788-37-5
(3,5-Dimethoxyphenyl)acetic acid	4670-10-4
2,3-Dimethoxyphenylacetic acid	
Manganie Nitrate	20694-39-7
Triphenylphosphine selenide	3878-44-2
3,5-Dichlorophenylhydrazine	39943-56-1
1-(3-Dimethylaminopropyl)-piperazine	877-96-3
N-Methyl-2-phenylacetamide	6830-82-6
Ethyl 3-methyl-4,4,4-trifluorobutyrate	95853-67-1
2-Naphthyl-beta-D-galactopyranoside	33993-25-8
2-Nitro-4-(trifluoromethyl)benzoic acid	320-94-5
3-Nitro-5-(trifluoromethyl)benzoic acid	328-80-3
2,2-Diethoxyacetamide	61189-99-9
Glyoxylic amide diethyl acetal	

3-(4-Methoxyphenyl)pyrazole

27069-17-6

3-Bromo-1,1,1-trifluoro-2-propanol

88378-50-1

431-34-5

Octadecylbenzene

4445-07-2

Stearylbenzene

1-Phenyloctadecane

ChemExper Chemical Directory : catalog of chemicals and suppliers

CHEMEXPER**MSDS online-Official Site**

Over 2 million MSDS documents Easy to use,
plus Free 30-day trial
www.msdsonline.com

Health & Safety Software

User configurable software for Environmental,
Health and Safety
www.etq.com

Chemical/Hazmat Software

Cradle to Grave Web App includes: MSDS, Tier
2, Compliance, much more
www.environmax.com

[Ads by Goooooogle](#)[Advertise on this site](#)

PRODUCT > NH

2,7-DIBROMO-9-FLUORENONE

PRODUCT IDENTIFICATION

CAS NO. 14348-75-5

EINECS NO.

FORMULA $C_{13}H_6Br_2O$

MOL WT. 337.99

H.S. CODE

TOXICITY

SYNONYMS 2,7-Dibromo Fluorene-9-one; 2,7-Dibromo Fluorenone;

DERIVATION

CLASSIFICATION

PHYSICAL AND CHEMICAL PROPERTIES

PHYSICAL STATE yellow crystalline powder

MELTING POINT 202 - 204 C

BOILING POINT

SPECIFIC GRAVITY

SOLUBILITY IN WATER

AUTOIGNITION

pH

VAPOR DENSITY

NFPA RATINGS

REFRACTIVE INDEX

FLASH POINT

STABILITY Stable under ordinary conditions. Oxidizes in light

APPLICATIONS

Fluorene is a member of polycyclic aromatic hydrocarbon (PAH). Two benzene rings are fused to cyclopentane ring. It emits violet fluorescent color. It is not synthesized commercially but is obtained from middle oil fraction of coal tar. It is insoluble in water; soluble in ether and acetone; melting point 116-117 C. It plays important part in metallocene catalysts as a ligand. It is used in the formation of polyradicals for resins. It is used in manufacturing antimalaria drugs and other pharmaceuticals. Fluorene family compounds are base materials for dyes and optical brightening agents. They have useful functions such as light and temperature sensitivity, heat resistance, conductivity, emittability, corrosion resistance and detection of amino groups. They are used in the applications of thermo and light sensitizer, liquid crystal chemistry, luminescence chemistry, spectrophotometric analysis, molecular chemistry, organometallic-complexes and biochemorphology industry. Oxidation of fluorene occur readily at the most reactive position 9. 9-Fluorenone has been investigated as an attractive element in organic solar cells, and display devices.

SALES SPECIFICATION

APPEARANCE yellow crystalline powder

ASSAY (G.C) 97.0% min

MELTING POINT 202 - 204 C


TRANSPORTATION

PACKING 20kgs in fiber drum

HAZARD CLASS

UN NO.

OTHER INFORMATION



European Hazard Symbols: XI, Risk Phrases: 36/37/38, Safety Phrases: 26, 37/39

We are pleased to receive your any questions or remarks. You may contact us at the following numbers and e-mail address for additional information.

Fax : + 82 + 2 + 783-8063 , + 82 + 2 + 3775-3073 (Seoul, Korea)

[HOME](#)[E-MAIL](#)